

RESEARCH PAPER RP1650

Part of *Journal of Research of the National Bureau of Standards*, Volume 34,
April 1945

HEATS AND FREE ENERGIES OF FORMATION OF THE PARAFFIN HYDROCARBONS, IN THE GASEOUS STATE, TO 1,500° K¹

By Edward J. Prosen, Kenneth S. Pitzer,² and Frederick D. Rossini

ABSTRACT

Values are presented for the heats of formation and the free energies of formation, from solid carbon (graphite) and gaseous hydrogen, of all the normal paraffin hydrocarbons and of the isomeric paraffins from the butanes through the octanes, in the gaseous state, to 1,500° K.

CONTENTS

	Page
I. Introduction.....	403
II. Unit of energy and molecular weights.....	403
III. Method of calculation.....	404
IV. Data used in the calculations.....	404
V. Results of the calculations.....	405
VI. References.....	411

I. INTRODUCTION

This paper presents new values for the heats and free energies of formation, from solid carbon (graphite) and gaseous hydrogen, of each of the 40 paraffin hydrocarbons from methane through the octanes, and of the higher normal paraffins, in the gaseous state, to 1,500° K.

II. UNIT OF ENERGY AND MOLECULAR WEIGHTS

The unit of energy used in this paper is the conventional thermochemical calorie defined as follows [1].³

1 calorie=4.1833 international (NBS) joules.

The molecular weights were calculated from the values given in the 1941 table of International Atomic Weights [2].

¹ This investigation was performed at the National Bureau of Standards jointly by the Thermochemical Laboratory and the American Petroleum Institute Research Project 44 on the "Collection and Analysis of Data on the Properties of Hydrocarbons."

² Consultant to the American Petroleum Institute Research Project 44 Professor of Chemistry, University of California, Berkeley, Calif.

³ Figures in brackets indicate the literature references at the end of this paper.

III. METHOD OF CALCULATION

Values of ΔH_f° , the increment in heat content for the reaction of forming the given hydrocarbon in the gaseous state, from the elements carbon (solid, graphite) and hydrogen (gas), with the reactants and the product in their appropriate thermodynamic standard states, at 0°K , were calculated by combining values of $\Delta H_f^\circ_{298.16}$ with values of $H^\circ_{298.16} - H^\circ_0$ for the appropriate compounds, by means of the relation ⁴

$$\Delta H_f^\circ = \Delta H_f^\circ_{298.16} - \Delta(H^\circ_{298.16} - H^\circ_0), \quad (1)$$

where the last term on the right is the value of $H^\circ_{298.16} - H^\circ_0$ (the heat content at 25°C less that at 0°K) for 1 mole of the gaseous hydrocarbon less that of n moles of solid carbon (graphite) and $(n+1)$ moles of gaseous hydrogen (the molecular formula of the hydrocarbon being $\text{C}_n\text{H}_{2n+2}$).

Values of ΔH_f° for the other temperatures were then systematically calculated by means of the relation

$$\Delta H_f^\circ = \Delta H_f^\circ + \Delta(H^\circ - H^\circ_0). \quad (2)$$

Values of ΔF_f° , the increment in free energy for the reaction of forming the given hydrocarbon in the gaseous state, from the elements carbon (solid, graphite) and hydrogen (gas), with the reactants and the product in their appropriate thermodynamic standard states at the given temperature, were calculated by combining values of ΔH_f° with appropriate values of $(F^\circ - H^\circ_0)/T$, the free-energy function, by means of the relation

$$\Delta F_f^\circ = \Delta H_f^\circ + T\Delta(F^\circ - H^\circ_0)/T, \quad (3)$$

where the last term on the right is the value of the product of the absolute temperature and $\Delta(F^\circ - H^\circ_0)/T$, the free energy at the given temperature less the heat content at 0°K , divided by the absolute temperature, for 1 mole of the gaseous hydrocarbon less that of n moles of solid carbon (graphite) and $(n+1)$ moles of gaseous hydrogen (the molecular formula of the hydrocarbon being $\text{C}_n\text{H}_{2n+2}$).

IV. DATA USED IN THE CALCULATIONS

The data used in the calculations include the following:

Values of the heat of formation, ΔH_f° , for the gaseous hydrocarbons at 25°C are taken from reference [4].

Values of the heat content at a given temperature less that at 0°K , $H^\circ - H^\circ_0$, and of the free-energy function, $(F^\circ - H^\circ_0)/T$, for the gaseous hydrocarbons, are taken from the following sources: For methane, from reference [5]; for ethane, propane, *n*-butane, *n*-pentane, *n*-hexane, *n*-heptane, *n*-octane, and the higher normal paraffins, from refer-

⁴ The usual convention is followed of letting the subscript denote the absolute temperature. The superscript denotes the thermodynamic standard state. See reference [3].

ence [6]; for isobutane, isopentane, and neopentane, from reference [7]; for the difference between *n*-hexane and its isomers and between *n*-heptane and its isomers, from reference [8]; for the isomeric octanes, from references [7] and [9]. Values of the foregoing properties for several of the intermediate temperatures were calculated by W. J. Taylor of this Bureau in connection with the work of the American Petroleum Institute Research Project 44.

V. RESULTS OF THE CALCULATIONS

The resulting values ⁵ of the heat of formation, ΔH_f° , are given in table 1. The resulting values ⁵ of the free energy of formation, ΔF_f° , are given in table 2.

⁵ The uncertainty in the values for 2,3-dimethylhexane may be several times that for the other isomeric octanes [4].

TABLE 1.—Values of the heats of formation of the paraffin hydrocarbons in the gaseous state to 1,500° K

Compound (gas)	Temperature in °K														
	0	298.16	300	400	500	600	700	800	900	1,000	1,100	1,200	1,300	1,400	1,500
	ΔH_f° in kcal/mole														
Methane.....	-15.987	-17.889	-17.903	-18.629	-19.302	-19.893	-20.401	-20.823	-21.166	-21.43	-21.65	-21.79	-21.92	-22.00	-22.06
Ethane.....	-16.517	-20.236	-20.258	-21.419	-22.437	-23.28	-23.98	-24.53	-24.97	-25.28	-25.50	-25.64	-25.72	-25.75	-25.73
Propane.....	-19.482	-24.820	-24.848	-26.358	-27.622	-28.66	-29.49	-30.12	-30.58	-30.90	-31.09	-31.16	-31.19	-31.14	-31.06
<i>n</i> -Butane.....	-23.332	-29.812	-29.847	-31.629	-33.128	-34.33	-35.28	-36.00	-36.54	-36.88	-37.06	-37.11	-37.05	-36.95	-36.79
2-Methylpropane.....	-24.602	-31.452	-31.489	-33.303	-34.783	-35.97	-36.91	-37.61	-38.10	-38.42	-38.57	-38.58	-38.54	-38.43	-38.28
<i>n</i> -Pentane.....	-27.27	-35.00	-35.04	-37.12	-38.86	-40.25	-41.34	-42.16	-42.75	-43.10	-43.26	-43.27	-43.18	-42.99	-42.74
2-Methylbutane.....	-28.66	-36.92	-36.96	-39.07	-40.78	-42.12	-43.17	-43.94	-44.47	-44.80	-44.92	-44.88	-44.78	-44.58	-44.36
2,2-Dimethylpropane.....	-31.30	-39.67	-39.71	-41.77	-43.42	-44.67	-45.63	-46.29	-46.73	-46.96	-47.01	-46.90	-46.70	-46.42	-46.12
<i>n</i> -Hexane.....	-30.98	-39.96	-40.01	-42.38	-44.36	-45.93	-47.15	-48.06	-48.71	-49.07	-49.23	-49.19	-49.04	-48.78	-48.47
2-Methylpentane.....	-32.03	-41.66	-41.71	-44.04	-45.92	-47.35	-48.43	-49.23	-49.84	-50.27					
3-Methylpentane.....	-31.36	-41.02	-41.07	-43.40	-45.27	-46.68	-47.73	-48.49	-49.05	-49.40					
2,2-Dimethylbutane.....	-34.52	-44.35	-44.40	-46.76	-48.64	-50.03	-51.05	-51.75	-52.22	-52.46					
2,3-Dimethylbutane.....	-32.82	-42.49	-42.54	-44.81	-46.59	-47.93	-48.92	-49.65	-50.23	-50.65					
<i>n</i> -Heptane.....	-34.65	-44.89	-44.94	-47.60	-49.83	-51.58	-52.93	-53.94	-54.63	-55.01	-55.15	-55.08	-54.88	-54.55	-54.16
2-Methylhexane.....	-35.77	-46.60	-46.66	-49.30	-51.45	-53.09	-54.33	-55.25	-55.91	-56.33					
3-Methylhexane.....	-34.96	-45.96	-46.02	-48.71	-50.90	-52.58	-53.84	-54.77	-55.42	-55.82					
3-Ethylpentane.....	-34.10	-45.34	-45.40	-48.15	-50.40	-52.12	-53.43	-54.41	-55.11	-55.56					
2,2-Dimethylpentane.....	-38.00	-49.29	-49.35	-52.02	-54.14	-55.72	-56.87	-57.69	-58.25	-58.56					
2,3-Dimethylpentane.....	-36.29	-47.62	-47.68	-50.39	-52.56	-54.21	-55.43	-56.35	-57.01	-57.45					
2,4-Dimethylpentane.....	-36.98	-48.30	-48.36	-51.02	-53.13	-54.70	-55.85	-56.69	-57.27	-57.64					
3,3-Dimethylpentane.....	-36.92	-48.17	-48.23	-50.88	-52.98	-54.54	-55.67	-56.46	-56.98	-57.28					
2,2,3-Trimethylbutane.....	-37.57	-48.96	-49.02	-51.64	-53.69	-55.19	-56.25	-57.01	-57.52	-57.83					
<i>n</i> -Octane.....	-38.33	-49.82	-49.88	-52.83	-55.30	-57.23	-58.71	-59.81	-60.67	-60.96	-61.08	-60.97	-60.71	-60.31	-59.86
2-Methylheptane.....	-39.42	-51.50	-51.56	-54.49	-56.87	-58.67	-60.04	-61.13	-61.91	-62.19					
3-Methylheptane.....	-38.64	-50.82	-50.88	-53.84	-56.23	-58.03	-59.40	-60.49	-61.27	-61.57					
4-Methylheptane.....	-38.43	-50.69	-50.75	-53.73	-56.13	-57.97	-59.39	-60.47	-61.23	-61.55					
3-Ethylhexane.....	-37.71	-50.40	-50.46	-53.59	-56.10	-58.04	-59.53	-60.64	-61.42	-61.81					
2,2-Dimethylhexane.....	-41.23	-53.71	-53.77	-56.72	-59.06	-60.85	-62.16	-63.02	-63.66	-63.96					
2,3-Dimethylhexane.....	-38.76	-51.13	-51.19	-54.08	-56.26	-57.88	-59.11	-60.10	-60.83	-61.16					
2,4-Dimethylhexane.....	-39.74	-52.44	-52.51	-55.50	-57.92	-59.72	-61.06	-62.10	-62.86	-63.23					
2,5-Dimethylhexane.....	-40.61	-53.21	-53.27	-56.24	-58.58	-60.33	-61.66	-62.70	-63.45	-63.77					
3,3-Dimethylhexane.....	-39.90	-52.61	-52.68	-55.65	-58.03	-59.78	-61.04	-61.93	-62.54	-62.84					

3,4-Dimethylhexane.....	-38.52	-50.91	-50.97	-53.81	-56.05	-57.71	-58.98	-60.05	-60.82	-61.01	-----	-----	-----	-----	
2-Methyl-3-ethylpentane.....	-37.96	-50.48	-50.54	-53.46	-55.77	-57.52	-58.87	-59.95	-60.70	-60.93	-----	-----	-----	-----	
3-Methyl-3-ethylpentane.....	-38.68	-51.38	-51.45	-54.32	-56.65	-58.42	-59.74	-60.65	-61.18	-61.29	-----	-----	-----	-----	
2,2,3-Trimethylpentane.....	-39.77	-52.61	-52.68	-55.62	-57.92	-59.63	-60.89	-61.82	-62.43	-62.63	-----	-----	-----	-----	
2,2,4-Trimethylpentane.....	-40.73	-53.57	-53.64	-56.58	-58.88	-60.59	-61.85	-62.78	-63.39	-63.59	-----	-----	-----	-----	
2,3,3-Trimethylpentane.....	-39.01	-51.73	-51.80	-54.72	-56.98	-58.66	-59.93	-60.92	-61.56	-61.62	-----	-----	-----	-----	
2,3,4-Trimethylpentane.....	-39.12	-51.97	-52.04	-54.97	-57.18	-58.83	-60.11	-61.17	-61.91	-62.03	-----	-----	-----	-----	
2,2,3,3-Tetramethylbutane.....	-41.09	-53.99	-54.06	-56.96	-59.14	-60.65	-61.70	-62.51	-63.02	-63.03	-----	-----	-----	-----	
<i>n</i> -Nonane.....	-42.00	-54.74	-54.81	-58.05	-60.76	-62.88	-64.49	-65.68	-66.50	-66.90	-67.01	-66.85	-66.54	-66.07	-65.55
<i>n</i> -Decane.....	-45.67	-59.67	-59.74	-63.28	-66.23	-68.53	-70.27	-71.55	-72.42	-72.84	-72.94	-72.74	-72.37	-71.82	-71.24
<i>n</i> -Undecane.....	-49.34	-65.60	-64.67	-68.50	-71.69	-74.17	-76.04	-77.42	-78.35	-78.78	-78.87	-78.63	-78.20	-77.58	-76.93
<i>n</i> -Dodecane.....	-53.02	-69.52	-69.60	-73.72	-77.16	-79.82	-81.82	-83.29	-84.28	-84.72	-84.79	-84.51	-84.03	-83.34	-82.63
<i>n</i> -Tridecane.....	-56.69	-74.45	-74.53	-78.94	-82.62	-85.47	-87.60	-89.17	-90.20	-90.66	-90.72	-90.40	-89.86	-89.10	-88.32
<i>n</i> -Tetradecane.....	-60.36	-79.38	-79.46	-84.17	-88.09	-91.12	-93.38	-95.04	-96.13	-96.60	-96.65	-96.28	-95.69	-94.86	-94.01
<i>n</i> -Pentadecane.....	-64.04	-84.31	-84.39	-89.39	-93.55	-96.77	-99.16	-100.91	-102.06	-102.54	-102.57	-102.17	-101.52	-100.61	-99.70
<i>n</i> -Hexadecane.....	-67.71	-89.23	-89.33	-94.61	-99.02	-102.41	-104.93	-106.78	-107.99	-108.48	-108.50	-108.06	-107.35	-106.37	-105.39
<i>n</i> -Heptadecane.....	-71.38	-94.15	-94.26	-99.84	-104.48	-108.06	-110.71	-112.65	-113.91	-114.42	-114.43	-113.94	-113.19	-112.13	-111.08
<i>n</i> -Octadecane.....	-75.06	-99.08	-99.19	-105.06	-109.95	-113.71	-116.49	-118.52	-119.84	-120.37	-120.35	-119.83	-119.02	-117.89	-116.78
<i>n</i> -Nonadecane.....	-78.73	-104.00	-104.12	-110.28	-115.41	-119.36	-122.27	-124.39	-125.77	-126.31	-126.28	-125.71	-124.85	-123.64	-122.47
<i>n</i> -Eicosane.....	-82.40	-108.93	-109.05	-115.50	-120.88	-125.01	-128.04	-130.26	-131.69	-132.25	-132.21	-131.60	-130.68	-129.40	-128.16
Δ per CH ₂	-3.673	-4.926	-4.931	-5.223	-5.465	-5.648	-5.778	-5.871	-5.927	-5.941	-5.927	-5.886	-5.830	-5.758	-5.692

TABLE 2.—Values of the free energies of formation of the paraffin hydrocarbons in the gaseous state to 1,500 °K

Compound (gas)	Temperature in °K														
	0	298.16	300	400	500	600	700	800	900	1,000	1,100	1,200	1,300	1,400	1,500
	ΔF_f° in kcal/mole														
Methane.....	-15.987	-12.140	-12.105	-10.048	-7.841	-5.49	-3.05	-0.55	2.01	4.61	7.22	9.85	12.50	15.14	17.79
Ethane.....	-16.517	-7.860	-7.785	-3.447	1.168	5.97	10.90	15.92	21.00	26.13	31.28	36.45	41.62	46.79	51.99
Propane.....	-19.482	-5.614	-5.541	1.191	8.230	15.50	22.93	30.45	38.05	45.68	53.34	61.01	68.68	76.35	84.08
<i>n</i> -Butane.....	-23.332	-3.754	-3.595	5.435	14.868	24.59	34.50	44.50	54.59	64.71	74.87	85.05	95.25	105.42	115.61
2-Methylpropane.....	-24.602	-4.296	-4.127	5.273	15.088	25.18	35.44	45.82	56.29	66.79	77.32	87.86	98.39	108.90	119.44
<i>n</i> -Pentane.....	-27.27	-1.96	-1.76	9.66	21.57	33.79	46.22	58.77	71.41	84.10	96.81	109.55	122.30	135.01	147.75
2-Methylbutane.....	-28.66	-3.50	-3.38	8.26	20.29	32.66	45.19	57.84	70.62	83.41	96.23	109.07	121.89	134.68	147.53
2,2-Dimethylbutane.....	-31.30	-3.64	-3.42	8.99	21.89	35.08	48.46	61.93	75.49	89.08	102.68	116.27	129.88	143.44	157.00
<i>n</i> -Hexane.....	-30.98	0.05	0.30	14.11	28.46	43.18	58.13	73.21	88.40	103.64	118.90	134.19	149.48	164.75	180.04
2-Methylpentane.....	-32.03	-0.96	-0.71	13.25	27.79	42.72	57.88	73.15	88.51	103.86	-----	-----	-----	-----	-----
3-Methylpentane.....	-31.36	-0.29	-0.04	13.94	28.50	43.45	58.62	73.89	89.23	104.53	-----	-----	-----	-----	-----
2,2-Dimethylbutane.....	-34.52	-2.35	-2.09	12.30	27.30	42.69	58.29	73.97	89.71	105.37	-----	-----	-----	-----	-----
2,3-Dimethylbutane.....	-32.82	-0.73	-0.47	13.81	28.67	43.92	59.38	74.94	90.57	106.18	-----	-----	-----	-----	-----
<i>n</i> -Heptane.....	-34.65	2.09	2.38	18.58	35.39	52.60	70.08	87.70	105.43	123.22	141.02	158.85	176.69	194.50	212.38
2-Methylhexane.....	-35.77	0.98	1.27	17.61	34.60	52.01	69.67	87.47	105.33	123.18	-----	-----	-----	-----	-----
3-Methylhexane.....	-34.96	1.10	1.39	17.67	34.40	51.68	69.21	86.87	104.59	122.29	-----	-----	-----	-----	-----
3-Ethylpentane.....	-34.10	2.59	2.88	19.36	36.52	54.15	72.06	90.09	108.19	126.25	-----	-----	-----	-----	-----
2,2-Dimethylpentane.....	-38.00	0.09	0.39	17.33	34.95	53.01	71.33	89.73	108.18	126.55	-----	-----	-----	-----	-----
2,3-Dimethylpentane.....	-36.29	0.16	0.45	16.83	33.90	51.43	69.24	87.17	105.18	123.16	-----	-----	-----	-----	-----
2,4-Dimethylpentane.....	-36.98	0.72	1.02	17.81	35.28	53.19	71.37	89.66	107.99	126.27	-----	-----	-----	-----	-----
3,3-Dimethylpentane.....	-36.92	0.63	0.93	17.64	35.02	52.84	70.92	89.10	107.34	125.53	-----	-----	-----	-----	-----
2,2,3-Trimethylbutane.....	-37.57	0.76	1.06	18.06	35.75	53.87	72.24	90.72	109.23	127.68	-----	-----	-----	-----	-----
<i>n</i> -Octane.....	-38.33	4.14	4.47	23.06	42.32	62.03	82.03	102.17	122.45	142.79	163.14	183.51	203.90	224.26	244.71
2-Methylheptane.....	-39.42	3.06	3.39	22.14	41.58	61.47	81.63	101.88	122.27	142.83	-----	-----	-----	-----	-----
3-Methylheptane.....	-38.64	3.29	3.62	22.21	41.50	61.27	81.31	101.44	121.68	142.07	-----	-----	-----	-----	-----
4-Methylheptane.....	-38.43	4.00	4.34	23.12	42.62	62.61	82.88	103.23	123.68	144.26	-----	-----	-----	-----	-----
3-Ethylhexane.....	-37.71	3.95	4.28	22.94	42.39	62.37	82.64	102.98	123.42	143.98	-----	-----	-----	-----	-----
2,2-Dimethylhexane.....	-41.23	2.56	2.91	22.18	42.20	62.71	83.48	104.30	125.21	146.27	-----	-----	-----	-----	-----

2,3-Dimethylhexane	-38.76	4.23	4.57	23.51	43.16	63.27	83.64	104.09	124.63	145.30					
2,4-Dimethylhexane	-39.74	2.80	3.14	22.05	41.72	61.89	82.32	102.84	123.43	144.07					
2,5-Dimethylhexane	-40.61	2.50	2.84	21.92	41.75	62.08	82.69	103.41	124.20	144.99					
3,3-Dimethylhexane	-39.90	3.17	3.51	22.63	42.50	62.85	83.47	104.16	124.93	145.76					
3,4-Dimethylhexane	-38.52	4.97	5.31	24.42	44.27	64.57	85.12	105.75	126.49	147.35					
2-Methyl-3-ethylpentane	-37.96	5.08	5.43	24.47	44.24	64.47	84.96	105.52	126.21	147.09					
3-Methyl-3-ethylpentane	-38.68	4.76	5.11	24.32	44.30	64.77	85.48	106.22	127.05	148.01					
2,2,3-Trimethylpentane	-39.77	4.09	4.44	23.82	43.98	64.63	85.54	106.52	127.57	148.68					
2,2,4-Trimethylpentane	-40.73	3.13	3.48	22.86	43.02	63.67	84.58	105.56	126.61	147.72					
2,3,3-Trimethylpentane	-39.01	4.52	4.86	24.09	44.08	64.55	85.30	106.10	126.98	147.90					
2,3,4-Trimethylpentane	-39.12	4.32	4.67	23.89	43.89	64.37	85.11	105.94	126.84	147.80					
2,2,3,3-Tetramethylbutane	-41.09	4.88	5.24	25.34	46.19	67.47	88.97	110.52	132.19	154.05					
<i>n</i> -Nonane	-42.00	6.18	6.56	27.54	49.25	71.46	93.98	116.66	139.48	162.37	184.20	208.18	231.12	254.03	277.04
<i>n</i> -Decane	-45.67	8.23	8.65	32.01	56.18	80.89	105.93	131.14	156.51	181.95	205.26	232.85	258.33	283.79	309.37
<i>n</i> -Undecane	-49.34	10.28	10.74	36.49	63.11	90.31	117.88	145.63	173.54	201.63	226.32	257.52	285.54	313.55	341.70
<i>n</i> -Dodecane	-53.02	12.33	12.83	40.97	70.04	99.74	129.83	160.11	190.57	221.11	247.38	282.19	312.76	343.32	374.03
<i>n</i> -Tridecane	-56.69	14.37	14.92	45.45	76.97	109.17	141.78	174.59	207.60	240.69	268.45	306.86	339.97	373.08	406.36
<i>n</i> -Tetradecane	-60.36	16.42	17.01	49.93	83.90	118.60	153.73	189.08	224.63	260.26	289.51	331.52	367.19	402.85	438.69
<i>n</i> -Pentadecane	-64.04	18.47	19.10	54.41	90.83	128.03	165.68	203.56	241.66	279.84	310.57	356.19	394.40	432.61	471.02
<i>n</i> -Hexadecane	-67.71	20.52	21.19	58.89	97.76	137.46	177.63	218.04	258.69	299.42	331.63	380.86	421.62	462.38	503.36
<i>n</i> -Heptadecane	-71.38	22.56	23.27	63.36	104.69	146.88	189.58	232.53	275.72	319.00	352.69	405.53	448.83	492.14	535.69
<i>n</i> -Octadecane	-75.06	24.61	25.36	67.84	111.62	156.31	201.53	247.01	292.75	338.58	373.75	430.20	476.05	521.90	568.02
<i>n</i> -Nonadecane	-78.73	26.66	27.45	72.32	118.55	165.74	213.48	261.49	309.78	358.16	394.81	454.86	503.26	551.67	600.35
<i>n</i> -Eicosane	-82.40	28.71	29.54	76.80	125.48	175.17	225.43	275.98	326.81	377.74	415.87	479.53	530.48	581.43	632.68
Δ per CH ₂	-3.673	2.048	2.090	4.479	6.931	9.428	11.951	14.484	17.029	19.579	21.061	24.668	27.215	29.764	32.331

Figure 1 is a plot that shows the thermodynamic stability, per carbon atom and with respect to the elements solid carbon (graphite) and gaseous hydrogen, of the normal paraffin hydrocarbons in the gaseous state as a function of the temperature. This plot shows the thermodynamic stability, per carbon atom, with respect to the ele-




FIGURE 1.—Thermodynamic stability of the normal paraffin hydrocarbons in the gaseous state as a function of temperature.

The scale of ordinates gives the value of $(1/n)(\Delta F^\circ/T)$ in calories per degree mole, where n is the number of carbon atoms per molecule, T is the absolute temperature in degrees Kelvin, and ΔF° is the standard free energy of formation of the hydrocarbon from the elements, solid carbon (graphite) and gaseous hydrogen, as given in table 2. The scale of abscissas gives the temperature in degrees Kelvin.

ments, of the normal paraffin hydrocarbons in the gaseous state. Points below the zero line indicate that the gaseous hydrocarbon in its standard state has a thermodynamic tendency to be formed from solid carbon (graphite) and gaseous hydrogen in their respective standard states.

Plots showing the relative stability of *n*-butane and isobutane, the 3 pentanes, the 5 hexanes, the 9 heptanes, and the 18 octanes, are given in references [9 and 10].

VI. REFERENCES

- [1] E. F. Mueller and F. D. Rossini, *Am. J. Physics* **12**, 1 (1944).
- [2] G. P. Baxter, M. Guichard, O. Hönigschmid, and R. Whytlaw-Gray, *J. Am. Chem. Soc.* **63**, 845 (1941).
- [3] G. N. Lewis and M. Randall, *Thermodynamics and the Free Energy of Chemical Substances* (McGraw-Hill Book Co., Inc., New York, N. Y., 1923).
- [4] E. J. Prosen and F. D. Rossini, *J. Research NBS* **34**, 263 (1945) RP1642.
- [5] D. D. Wagman, J. E. Kilpatrick, W. J. Taylor, K. S. Pitzer, and F. D. Rossini, *J. Research NBS* **34**, 143 (1945) RP1634.
- [6] K. S. Pitzer, *Ind. Eng. Chem.* **36**, 829 (1944).
- [7] K. S. Pitzer and J. E. Kilpatrick, National Bureau of Standards. Unpublished data.
- [8] K. S. Pitzer. *Chem. Rev.* **27**, 39 (1940).
- [9] E. J. Prosen, K. S. Pitzer, and F. D. Rossini, *J. Research NBS* **34**, 255 (1945) RP1641.
- [10] F. D. Rossini, E. J. Prosen, and K. S. Pitzer, *J. Research NBS* **27**, 529 (1941) RP1440.

WASHINGTON, SEPTEMBER 15, 1944.



NATIONAL BUREAU OF STANDARDS, WASHINGTON 25, D. C.

Send me the Mathematical Tables marked X below. I enclose remittance to cover the cost.

Mark X	Title of publication	United States and its possessions, and countries extending franking privilege	Other countries	Amount enclosed
-----	MT2. Tables of the exponential function e^x -----	\$2. 00	\$2. 50	-----
-----	MT3. Tables of circular and hyperbolic sines and cosines for radian arguments-----	2. 00	2. 50	-----
-----	MT4. Tables of sines and cosines for radian arguments-----	2. 00	2. 50	-----
-----	MT5. Tables of sine, cosine, and exponential integrals, volume I-----	2. 00	2. 50	-----
-----	MT6. Tables of sine, cosine, and exponential integrals, volume II-----	2. 00	2. 50	-----
-----	MT7. Table of natural logarithms, volume I (1 to 50,000)-----	2. 00	2. 50	-----
-----	MT8. Tables of probability functions, volume I-----	2. 00	2. 50	-----
-----	MT9. Table of natural logarithms, volume II (50,000 to 100,000)-----	2. 00	2. 50	-----
-----	MT10. Table of natural logarithms, volume III (0.0001 to 5.0000)-----	2. 00	2. 50	-----
-----	MT11. Tables of moments of inertia and section moduli of ordinary angles, channels, and bulb angles with certain plate combinations-----	2. 00	2. 50	-----
-----	MT12. Table of natural logarithms, volume IV (5.0000 to 10.0000)-----	2. 00	2. 50	-----
-----	MT13. Table of sine and cosine integrals for arguments from 10 to 100-----	2. 00	2. 50	-----
-----	MT14. Tables of probability functions, volume II-----	2. 00	2. 50	-----
-----	MT15. The hypergeometric and Legendre functions with applications to integral equations of potential theory-----	2. 00	2. 50	-----
-----	MT16. Table of arc tan x -----	2. 00	2. 50	-----
-----	MT17. Miscellaneous physical tables—Planck's radiation functions: electronic functions-----	1. 50	1. 75	-----
-----	MT18. Table of the zeros of the Legendre polynomials of order 1-16 and the weight coefficients for Gauss' mechanical quadrature formula-----	. 25	. 30	-----
-----	MT19. On the function $H(m, a, x) = \exp(-ix)F(m+1-ia, 2m+2; ix)$ -----	. 25	. 30	-----
-----	MT20. Table of integrals $\int_0^x J_0(t) dt$ and $\int_0^x Y_0(t) dt$ -----	. 25	. 30	-----
-----	MT21. Table of $Ji_0(x) = \int_x^\infty \frac{J_0(t)}{t} dt$ and related functions-----	. 25	. 30	-----
-----	Carried forward to other side-----	-----	-----	-----

Mark X	Title of Publication	United States and its possessions, and countries ex- tending franking privilege	Other countries	Amount enclosed
-----	Brought forward from other side.....	-----	-----	-----
-----	MT22. Table of coefficients in numerical integration formulae.....	\$0. 25	\$0. 30	-----
-----	MT23. Table of Fourier coefficients.....	. 25	. 30	-----
-----	MT24. Coefficients for numerical differentiation with central differences.....	. 25	. 30	-----
-----	MT25. Seven-point Lagrangian integration formulas.....	. 25	. 30	-----
-----	MT26. A short table of the first five zeros of the transcendental equation, $J_0(x)Y_0(kx) - J_0(kx)Y_0(x) = 0$ 25	. 30	-----
-----	MT27. Table of coefficients for inverse interpolation with central differences.....	. 25	. 30	-----
-----	MT28. Table of $f_n(x) = \frac{n!}{(x/2)^n} J_n(x)$ 25	. 30	-----
-----	MT29. Table of coefficients for inverse interpolation with advancing differences.....	. 25	. 30	-----
-----	MT30. A new formula for inverse interpolation.....	. 25	. 30	-----
-----	MT31. Coefficients for interpolation within a square grid in the complex plane.....	. 25	. 30	-----
-----	MT32. Table of coefficients for differences in terms of the derivatives.....	. 25	. 30	-----
-----	Total remittance.....	-----	-----	-----

Remittance should be in form of post-office money order, or check, and made payable to the order of the "National Bureau of Standards"
in United States currency

Send to.....

Number and Street..... City, Zone, and State.....

MATHEMATICAL TABLES

Attention is invited to a series of publications prepared by the *Project for the Computation of Mathematical Tables* conducted by the Federal Works Agency, Work Projects Administration for the City of New York, under the sponsorship of the National Bureau of Standards. The tables which have been made available through the National Bureau of Standards are listed below. A list of other WPA tables obtainable elsewhere will be sent by the Bureau on request.

There is included in this list a publication of the hypergeometric and Legendre functions (MT15), prepared by the Bureau.

MT1. TABLE OF THE FIRST TEN POWERS OF THE INTEGERS FROM 1 TO 1,000: (1938) VIII + 80 pages; heavy paper cover. Out of print.

MT2. TABLES OF THE EXPONENTIAL FUNCTION e^x : The ranges and intervals of the argument and the number of decimal places in the entries are given below:

Range of x	Interval of x	Decimals given
-2.5000 to 1.0000	0.0001	18
1.0000 to 2.5000	.0001	15
2.500 to 5.000	.001	15
5.00 to 10.00	.01	12

(1939) XV + 535 pages; bound in buckram, \$2.00.

MT3. TABLES OF CIRCULAR AND HYPERBOLIC SINES AND COSINES FOR RADIAN ARGUMENTS: Contains 9 decimal place values of $\sin x$, $\cos x$, $\sinh x$, and $\cosh x$ for x (in radians) ranging from 0 to 2 at intervals of 0.0001. (1939) XVII + 405 pages; bound in buckram, \$2.00.

MT4. TABLES OF SINES AND COSINES FOR RADIAN ARGUMENTS: Contains 8 decimal place values of sines and cosines for radian arguments ranging from 0 to 25 at intervals of 0.001. (1940) XXIX + 275 pages; bound in buckram, \$2.00.

MT5. TABLES OF SINE, COSINE, AND EXPONENTIAL INTEGRALS, VOLUME I: Values of these functions to 9 places of decimals from 0 to 2 at intervals of 0.0001. (1940) XXVI + 444 pages; bound in buckram, \$2.00.

MT6. TABLES OF SINE, COSINE, AND EXPONENTIAL INTEGRALS, VOLUME II: Values of these functions to 9, 10, or 11 significant figures from 0 to 10 at intervals of 0.001 with auxiliary tables. (1940) XXXVII + 225 pages; bound in buckram, \$2.00.

MT7. TABLE OF NATURAL LOGARITHMS, VOLUME I: Logarithms of the integers from 1 to 50,000 to 16 places of decimals. (1941) XVIII + 501 pages; bound in buckram, \$2.00.

MT8. TABLES OF PROBABILITY FUNCTIONS, VOLUME I: Values of these functions to 15 places of decimals from 0 to 1 at intervals of 0.0001 and from 1 to 5.6 at intervals of 0.001. (1941) XXVIII + 302 pages; bound in buckram, \$2.00.

MT9. TABLE OF NATURAL LOGARITHMS, VOLUME II: Logarithms of the integers from 50,000 to 100,000 to 16 places of decimals. (1941) XVIII + 501 pages; bound in buckram, \$2.00.

MT10. TABLE OF NATURAL LOGARITHMS, VOLUME III: Logarithms of the decimal numbers from 0.0001 to 5.0000, to 16 places of decimals. (1941) XVIII + 501 pages; bound in buckram, \$2.00.

MT11. TABLES OF THE MOMENTS OF INERTIA AND SECTION MODULI OF ORDINARY ANGLES, CHANNELS, AND BULB ANGLES WITH CERTAIN PLATE COMBINATIONS: (1941) XIII + 197 pages; bound in green cloth. \$2.00.

MT12. TABLE OF NATURAL LOGARITHMS, VOLUME IV: Logarithms of the decimal numbers from 5.0000 to 10.0000, to 16 places of decimals. (1941) XXII + 506 pages; bound in buckram, \$2.00.

MT13. TABLE OF SINE AND COSINE INTEGRALS FOR ARGUMENTS FROM 10 TO 100: (1942) XXXII + 185 pages; bound in buckram, \$2.00.

MT14. TABLES OF PROBABILITY FUNCTIONS, VOLUME II: Values of these functions to 15 places of decimals from 0 to 1 at intervals of 0.0001 and from 1 to 7.8 at intervals of 0.001. (1942) XXI + 344 pages; bound in buckram, \$2.00.

MT15. THE HYPERGEOMETRIC AND LEGENDRE FUNCTIONS WITH APPLICATIONS TO INTEGRAL EQUATIONS OF POTENTIAL THEORY. By Chester Snow, National Bureau of Standards. Reproduced from original handwritten manuscript. (1942) VII + 319 pages, bound in heavy paper cover. \$2.00.

MT16. TABLE OF ARC TAN x : Table of inverse tangents for positive values of the angle in radians. Second central differences are included for all entries. $x = [0(.001)7(.01)50(.1)300(1)2,000(10)10,000; 12D]$ (1942) XXV + 169 pages; bound in buckram, \$2.00.

[Continued on p. 4 of cover]

- MT17. MISCELLANEOUS PHYSICAL TABLES: PLANCK'S RADIATION FUNCTIONS (Originally published in the Journal of the Optical Society of America, February 1940); and ELECTRONIC FUNCTIONS (1941) VII+58 pages; bound in buckram, \$1.50.
- MT18. TABLE OF THE ZEROS OF THE LEGENDRE POLYNOMIALS OF ORDER 1-16 AND THE WEIGHT COEFFICIENTS FOR GAUSS' MECHANICAL QUADRATURE FORMULA. (Reprinted from Bul. Amer. Mathematical Society, October 1942.) 5 pages, with cover, 25 cents.
- MT19. ON THE FUNCTION $H(m, a, x) = \exp(-ix)F(m+1-ia, 2m+2; ix)$; with table of the confluent hypergeometric function and its first derivative. (Reprinted from J. Math. Phys., December 1942.) 20 pages, with cover, 25 cents.
- MT20. TABLE OF INTEGRALS $\int_0^x J_0(t)dt$ and $\int_0^x Y_0(t)dt$. (Reprinted from J. Math. Phys., May 1943.) 12 pages, with cover, 25 cents.
- MT21. TABLE OF $J_0(x) = \int_x^\infty \frac{J_0(t)}{t} dt$ AND RELATED FUNCTIONS. (Reprinted from J. Math. Phys., June 1943.) 7 pages, with cover, 25 cents.
- MT22. TABLE OF COEFFICIENTS IN NUMERICAL INTEGRATION FORMULAE. (Reprinted from J. Math. Phys., June 1943.) 2 pages, with cover, 25 cents.
- MT23. TABLE OF FOURIER COEFFICIENTS. (Reprinted from J. Math. Phys., Sept. 1943.) 11 pages, with cover, 25 cents.
- MT24. COEFFICIENTS FOR NUMERICAL DIFFERENTIATION WITH CENTRAL DIFFERENCES. (Reprinted from J. Math. Phys., Sept. 1943.) 21 pages, with cover, 25 cents.
- MT25. SEVEN-POINT LAGRANGIAN INTEGRATION FORMULAS. (Reprinted from J. Math. Phys., Dec. 1943.) 4 pages, with cover, 25 cents.
- MT26. A SHORT TABLE OF THE FIRST FIVE ZEROS OF THE TRANSCENDENTAL EQUATION. $J_0(x)Y_0(kx) - J_0(kx)Y_0(x) = 0$. (Reprinted from J. Math. Phys., Dec. 1943.) 2 pages, with cover, 25 cents.
- MT27. TABLE OF COEFFICIENTS FOR INVERSE INTERPOLATION WITH CENTRAL DIFFERENCES. (Reprinted from J. Math. Phys., Dec. 1943.) 15 pages, with cover, 25 cents.
- MT28. TABLE OF $f_n(x) = \frac{n!}{(x/2)^n} J_n(x)$. (Reprinted from J. Math. Phys., Feb. 1944) 16 pages, with cover, 25 cents.
- MT29. TABLE OF COEFFICIENTS FOR INVERSE INTERPOLATION WITH ADVANCING DIFFERENCES (Reprinted from J. Math. Phys., May 1944.) 28 pages, with cover, 25 cents.
- MT30. A NEW FORMULA FOR INVERSE INTERPOLATION. (Reprinted from Bul. Amer. Mathematical Society, Aug. 1944.) 4 pages, with cover, 25 cents.
- MT31. COEFFICIENTS FOR INTERPOLATION WITHIN A SQUARE GRID IN THE COMPLEX PLANE. (Reprinted from J. Math. Phys., Aug. 1944.) 11 pages, with cover. 25 cents.
- MT32. TABLE OF COEFFICIENTS FOR DIFFERENCES IN TERMS OF THE DERIVATIVES. (Reprinted from J. Math. Phys., Nov. 1944.) 4 pages, with cover, 25 cents.

Payment is required in advance. Make remittance payable to the "National Bureau of Standards" and send with order, using the blank form facing page 3 of the cover.

A mailing list is maintained for those who desire to receive announcements regarding new tables as they become available.